

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Rubbing behavior of European brown bears: Factors affecting rub tree selectivity and density

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1800262> since 2021-09-08T18:11:21Z

Published version:

DOI:10.1093/jmammal/gyaa170

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on:

Questa è la versione dell'autore dell'opera:

[Journal of Mammalogy, 2021, DOI: 10.1093/jmammal/gyaa170]

The definitive version is available at:

La versione definitiva è disponibile alla URL:

<https://academic.oup.com/jmammal/article/102/2/468/6123321>

Rubbing behavior of European brown bears: factors affecting rub tree selectivity and density

ENRIQUE GONZÁLEZ-BERNARDO^{1,2*}, CARLOTTA BAGNASCO³, GIULIA BOMBIERI^{1,4},
ALEJANDRA ZARZO-ARIAS¹, HÉCTOR RUIZ-VILLAR¹, ANA MORALES-GONZÁLEZ⁵, CINDY
LAMAMY⁶, ANDRÉS ORDIZ⁷, DAVID CAÑEDO⁸, JUAN DÍAZ⁸, DANIEL E. CHAMBERLAIN³,
AND VINCENZO PENTERIANI¹

¹*Research Unit of Biodiversity (UMIB, CSIC-UO-PA), Mieres Campus, 33600 Mieres, Spain (EGB, GB, AZA, HRV, VP)*

²*Pyrenean Institute of Ecology (IPE), C.S.I.C., Avda. Montañana 1005, 50059 Zaragoza, Spain (EGB)*

³*Dpt. of Life Sciences and Systems Biology, University of Turin, Via Accademia Albertina, 13-10123, Turin, Italy (CB, DEC)*

⁴*Museo delle Scienze, Sezione Zoologia dei Vertebrati, Corso del Lavoro e della Scienza 3, I-38123, Trento, Italy (GB)*

⁵*Estación Biológica de Doñana, C.S.I.C., Department of Conservation Biology, Avda. Americo Vespucio 26, 41092 Sevilla, Spain (AMG)*

⁶*Gembloux Agro-Bio Tech, Dpt. BIOSE, Liège University, Passage des Déportés, 5030, Gembloux, Belgium (CL)*

⁷*Faculty of Environmental Sciences and Natural Resource Management, Norwegian University of Life Sciences, Postbox 5003, NO-1432, Ås, Norway (AO)*

⁸*Consejería de Ordenación del Territorio, Infraestructuras y Medio Ambiente, Dirección General de Biodiversidad, Principado de Asturias, Oviedo, Spain (DC, JD)*

**Correspondent: gonzalezbernardoenrique@gmail.com*

Scent-mediated communication is considered the principal communication channel in many mammal species. Compared with visual and vocal communication, odors persist for a longer time, enabling individuals to interact without being in the same place at the same time. The brown bear (*Ursus arctos*), like other mammals, carries out chemical communication, for example, by means of scents deposited on marking (or rub) trees. In this study, we assessed rub tree selectivity of the brown bear in the predominantly deciduous forests of the Cantabrian Mountains (NW Spain). We first compared the characteristics of 101 brown bear rub trees with 263 control trees. We then analysed the potential factors affecting the density of rub trees along 35 survey routes along footpaths. We hypothesized that: (1) bears would select particular trees, or tree species, with characteristics that make them more conspicuous; and (2) that bears would select trees located in areas with the highest presence of conspecifics, depending on the population density or the position of the trees within the species' range. We used linear models and Generalized Additive Models to test these hypotheses. Our results showed that brown bears generally selected more conspicuous trees with a preference for birches (*Betula* spp.). This choice may facilitate the marking and/or detection of chemical signals and, therefore, the effectiveness of intraspecific communication. Conversely, the abundance of rub trees along footpaths did not seem to depend on the density of bear observations or their relative position within the population centre or its border. Our results suggest that Cantabrian brown bears select trees based on their individual characteristics and their location, with no influence of characteristics of the bear population itself. Our findings can be used to locate target trees that could help in population monitoring.

Keywords: *Ursus arctos*, rub trees, rubbing behaviour, tree selectivity, chemical communication

37 La comunicación olfativa se considera el principal canal de comunicación en muchas
38 especies de mamíferos. En comparación con la comunicación visual y la vocal, los
39 olores persisten durante más tiempo, lo que permite a los individuos interactuar sin estar
40 en el mismo lugar al mismo tiempo. El oso pardo (*Ursus arctos*), al igual que otros
41 mamíferos, emplea la comunicación química, por ejemplo, por medio de olores
42 depositados en árboles a través del marcaje o rascado. En este estudio, evaluamos la
43 selección de árboles de marcaje por el oso pardo en los bosques predominantemente
44 caducifolios de la Cordillera Cantábrica (noroeste de España). En primer lugar,
45 comparamos las características individuales de 101 árboles de marcaje de oso pardo con
46 263 árboles control. Después, analizamos los factores potenciales que afectan la
47 densidad de árboles de marcaje en 35 trayectos de prospección a lo largo de caminos y
48 pistas forestales. Planteamos las hipótesis que: (1) los osos seleccionan árboles
49 particulares, o especies de árboles, con características que los hacen más conspicuos; y
50 (2) que los osos seleccionan árboles ubicados en áreas con mayor presencia de
51 conespecíficos, dependiendo de la densidad de población osera o de la posición de los
52 árboles dentro del rango de distribución de la especie. Usamos modelos lineales y
53 modelos aditivos generalizados para probar estas hipótesis. Nuestros resultados
54 mostraron que los osos pardos generalmente seleccionaron árboles más conspicuos, con
55 preferencia por los abedules (*Betula* spp.). Esta elección puede facilitar el marcaje y/o
56 detección de señales químicas y, por tanto, la eficacia de la comunicación
57 intraespecífica. Por el contrario, la abundancia de marcaje a lo largo de los trayectos no
58 parece depender de la densidad de las observaciones de osos o de su posición relativa
59 con respecto al centro o los límites del rango de la población. Nuestros resultados
60 sugieren que los osos pardos cantábricos seleccionan árboles en función de sus
61 características individuales y de su ubicación, sin que influyan en ello las características

de la población osera. Nuestros hallazgos pueden servir para localizar árboles
específicos que podrían ayudar al monitoreo de la población.

Palabras clave: *Ursus arctos*, árboles de marcaje, comportamiento de marcaje,
comunicación química, selección de árboles

Chemical signals that are spread by distinct means, such as urine, feces, or glandular
secretions (Müller-Schwarze 2006; Johnston and Del Barco-Trillo 2010), are considered
the principal channel of communication in many mammal species (Ralls 1971), more so
than visual or acoustic signals (Müller-Schwarze 2006; Penteriani and Delgado 2017).
Mammalian scent-marking strategies mostly depend on individual and social factors, as
well as on the physical characteristics of the environment (Barja and De Miguel 2010).
Scent-mediated communication has several advantages, including persistence for long
periods, and facilitation of interactions among individuals without any need for direct
interaction at a given site (Mills et al. 1980).

Chemical communication can have different functions that include: self-
advertising, i.e., communication of the state or characteristics of the individual, such as
age, sex, reproductive status, and health condition; conspecific localisation;
communication of dominance; and the defence of a resource (Brown 1979; Doty 1986,
Gosling 1990; Johnston 2008; Morales-González et al. 2019). Sometimes, this form of
indirect interaction also can prevent agonistic encounters (Gosling and McKay 1990;
Roberts and Gosling 2001). While scent-marking plays an important role when
defending or signalling territory occupancy in territorial species (Roberts and Gosling
2001; Müller and Manser 2008; Barja and De Miguel 2010), scent marking in non-
territorial species is more difficult to interpret (Clapham et al. 2012). Solitary species

must maintain effective communication to sustain a social structure that facilitates reproduction (Clapperton 1989; Gosling and Roberts 2001), for example by communicating male fitness (White et al. 2002) or the reproductive status of females (Gorman and Trowbridge 1989; Rich and Hurst 1998; Barja and De Miguel 2010). In particular, large-bodied solitary carnivorous species with large home ranges base much of their communication on olfactory chemical signals (Macdonald 1980; Gorman and Trowbridge 1989; Lamb et al. 2017), which usually are left on conspicuous and permanent substrates, such as rocks and tree trunks (Kleiman 1966; Barja 2009; Allen et al. 2017). Thus, intensive marking behaviour on those substrates increases detectability and information transmission, reducing the investment of resources in communication (Alberts 1992; Gosling and Roberts 2001).

As in other solitary and non-territorial carnivores (Smith et al. 1989; Barja 2009), bears perform scent-marking on different substrates (Filipczyková et al. 2016). Indeed, rubbing behaviour has been documented in most ursid species, i.e., American black bear *Ursus americanus* (Burst and Pelton 1983; Sawaya et al. 2012) and Asiatic black bear *Ursus thibetanus* (Bromley 1965), giant panda *Ailuropoda melanoleuca* (White et al. 2002; Swaisgood et al. 2004; Nie et al. 2012), sloth bear *Melursus ursinus* (Laurie and Seidensticker 1977), Andean bear *Tremarctos ornatus* (Filipczyková et al. 2016; Kleiner et al. 2018) and brown bear *Ursus arctos* (Krott 1962; Shaffer 1971).

Brown bears exhibit a set of marking behaviours, including: urination; biting, scratching, and stripping bark; rubbing the back, shoulders, and head (Green and Mattson 2003; Puchkovskiy 2009; Clapham et al. 2012), and pedal and scent-marking at the base of trees and the surrounding area (Clapham et al. 2012; Sergiel et al. 2017). Marking, or rub trees (hereafter, RTs) have a key function in intraspecific communication in brown bears (Green and Mattson 2003; Clapham et al. 2012, 2013;

Sato et al. 2014; Tattoni et al. 2015; Lamb et al. 2017). Indeed, some studies have reported larger odoriferous glands and higher amounts of glandular secretion produced in individuals that show more pronounced rubbing behaviour (Tomiyasu et al. 2017). Other studies even have reported a different composition of glandular secretions between males and females (Rosell et al. 2011; Sergiel et al. 2017; Tomiyasu et al. 2017), which may facilitate identification of sex among bears (Jojola et al. 2012). In addition, the function of tree rubbing has been described as density-dependent, prioritizing the communication of dominance in situations of a high density of individuals (Lamb et al. 2017). There are two main functions and advantages of chemical scents on RTs: (a) communication of dominance or fitness in adult males (Clapham et al. 2012; Tattoni et al. 2015; Lamb et al. 2017) and related adult male avoidance by subadults (Jojola et al. 2012; Tomiyasu et al. 2017); and, (b) increasing the probability of finding a potential mate (Green and Mattson 2003; Clapham et al. 2012, 2014; Lamb et al. 2017).

The pool of RTs in a given area provides a communication network through which individuals exchange information; this pool can be used over time for generations (Green and Mattson 2003; Clapham et al. 2013; Morgan Henderson et al. 2015). RTs commonly are located in the proximity of foot trails or unpaved roads that facilitate the transit of bears (Lloyd 1979; Green and Mattson 2003; Sato et al. 2014), with trail-oriented rubbing marks (Green and Mattson 2003), or at forest edges (Green and Mattson 2003; Puchkovskiy 2009). Brown bears seem to select trees that allow for greater conspicuousness of their markings. Preference also is given to certain species of conifers (Puchkovskiy 2009; Sato et al. 2014), probably due to the resins that may enhance the detectability or range of scent-markings (Green and Mattson 2003,

Clapham et al. 2013; Tattoni et al. 2015). This effect has been suggested to be enhanced by producing wounds to the tree that increase resin flow (Sato et al. 2014).

Studies on RTs have mainly been focused on North American grizzlies *Ursus arctos horribilis* (Green and Mattson 2003; Clapham et al. 2013; Morgan Henderson et al. 2015), with a few also in Russia (Puchkovskiy 2009; Puchkovskiy et al. 2012) and Japan (Sato et al. 2014); however, these studies considered large continuous populations. To our knowledge, such a study has not previously been carried out in smaller, vulnerable populations, nor in populations living in areas dominated by deciduous forest. Here, by using (a) 101 RTs found in the Cantabrian Mountains (north-western Spain) and (b) the density of RTs along 35 survey routes along footpaths, we carried out a multilevel analysis to elucidate whether brown bears' RT selection is determined by tree species, physical traits, landscape position, and whether RT occurrence was related to their spatial location and observed bear density. Two main hypotheses underlie this study. First, as previously reported in populations inhabiting coniferous forests, RTs have characteristics that distinguish them from surrounding trees (e.g., dendrometric characteristics, tree location), which probably enhance the conspicuousness of chemical signalling and/or represent a better substrate for marking. We thus hypothesized that bears would select specific trees, or tree species with characteristics that make them more conspicuous. Second, we hypothesized that RTs would be more abundant in areas with the highest presence of conspecifics, towards the core of the bear distribution area and in areas where the density of bears is higher.

This study was carried out in a small and isolated, but still autochthonous, brown bear population, at the southwestern limit of the current species distribution. An overarching goal of the study therefore was to obtain information on scent-marking in small populations, whose behaviour may differ from larger and less isolated

populations, because the smaller population size is likely to mean a higher number of interactions between the same individuals. The results provide insights into the ecology of a threatened population and therefore contribute to conservation and monitoring actions.

METHODS

Study area.—The study area covers most of the geographic range of the western sub-population of brown bears in the Cantabrian Mountains (northwestern Spain, Figure 1), which run for ~300 km in parallel to the coast (E–W) with an average and maximum altitude of 1100 and 2648 m a.s.l, respectively (Martínez Cano et al. 2016; Penteriani et al. 2019). The region has an oceanic climate, more humid and temperate in northern slopes and continental and dryer on southern slopes (Ortega and Morales 2015). The landscape is covered predominantly by forests, shrubs, and croplands (Mateo-Sánchez et al. 2016). Forests of semi-deciduous and evergreen oaks (*Quercus* sp.) dominate sunny slopes, whereas the north-facing slopes are covered by deciduous trees such as beech (*Fagus sylvatica*) and common (*Q. robur*) and sessile oak (*Q. petraea*). Birches (*Betula* sp.) occupy areas of acid soils, dominant towards the west of the study area, in well illuminated areas with sufficient humidity and frequently colonizing degraded or bare areas, with few monospecific stands (García de Celis et al. 2004). Non-forested areas are mainly occupied by shrub species, such as heather (*Erica* sp., *Calluna* sp.) and brooms (*Cytisus* sp.; Fernández-Gil et al. 2006; Mateo-Sánchez et al. 2016). Above the treeline, berry-producing shrubs occur, including bilberries (*Vaccinium myrtillus*; Pato and Obeso 2012; Martínez Cano et al. 2016). Most of the areas inhabited by bears are surrounded by urbanized and cultivated areas, with a high density of transport

infrastructure; the main economic activities include livestock breeding, mining, timber harvesting, and recreational activities (Zarzo-Arias et al. 2018, 2019).

Description of the population.—In northern Spain, there are two brown bear populations that are isolated from other European populations, one in the Pyrenees and another in the Cantabrian Mountains, separated from each other by almost 300 km. The latter population is divided into two subpopulations, with little gene flow (Pérez et al. 2010; Gonzalez et al. 2016) and a genetic variability that is among the lowest of any brown bear population in the world (García-Garitagoitia et al. 2007). In the 1990s, the Cantabrian population consisted of a minimum of 70 – 85 bears (50 – 65 individuals in the western nucleus and 14 – 20 in the eastern; Clevenger and Purroy 1991; Naves and Palomero 1993; Pérez et al. 2014). Recent population trends seem positive, especially in the western subpopulation (Pérez et al. [2014] reported 168 – 260 bears in the western subpopulation and 12 – 40 in the eastern subpopulation), but this species is still considered as “Endangered” in Spain (BOE 2011).

Multilevel analysis

Location and characterisation of rub and control trees.—We surveyed principal trails and forest roads (hereafter footpaths) within the study area for RTs from October 2018 to March 2019 (Fig. 1). We looked for trees with rubbing signs such as smoothed bark, discoloured surface, scratches, bites, or lack of vegetation at the base; however, the tree was only considered an RT when the presence of bear fur snagged on the bark was confirmed (a characteristic sign of bear tree rubbing behaviour). All the trees located within a radius of 5 m around each marked tree (following Clapham et al. 2013) were considered control trees (hereafter, CT). None of the CTs were found to have any evidence of marking. This radius ensured that we sampled trees that showed the same local habitat characteristics as the RT, and that were clearly available to the bear in that

location. We followed previous studies on brown bear tree rubbing behaviour (Green and Mattson 2003; Clapham et al. 2013; Sato et al. 2014), to characterise each rub and control tree, recording the following variables: (1) tree species (categorical with five levels: birch [*Betula spp.*], oak [*Quercus spp.*], chestnut [*Castanea sativa*], conifer [*Pinus spp.*, *Pseudotsuga menziesii*], and other); (2) tree status (categorical with two levels: dead and alive); (3) other brown bear tree marks (categorical with three levels: bites, scratches, no other marks); (4) slope exposure, i.e., exposure where the tree was located (categorical with eight levels: north, northeast, east, southeast, south, southwest, west, northwest); (5) tree height, measured with NASA GLOBE Observer v 3.0 (NASA 2019); (6) trunk height, i.e., from the ground to the first branch, measured with a tape measure; (7) diameter at breast height (DBH), measured with a diameter tape; (8) distance to the nearest footpath (hereafter dist. to footpath), measured with a tape measure; (9) tree spacing, i.e., average distance to the nearest tree located in each of the four main cardinal directions, measured with a tape; and (10) terrain elevation (m.a.s.l.).

Characterization of the landscape around rub trees.—To explore the landscape characteristics of the areas in which the rub trees were found, both in terms of natural and human features of rub trees, we calculated the following variables for each rub tree: distance to the nearest (1) paved road (hereafter dist. to paved road); (2) unpaved road (hereafter dist. to unpaved road); (3) river (hereafter dist. to river); and (4) human settlement (hereafter dist. to human settlement). We attempted to identify the following potential landscape predictors of RTs by measuring each within a 1 km radius of CTs and RTs (based on Lamamy et al. 2019; Penteriani et al. 2020): (1) total paved road length; (2) total unpaved road length; (3) total river length; (4) ruggedness (i.e., total length of the linear development of level curves within the buffer); and (5) land use, expressed as the percentage occupied by each land cover class within the 1 km radius.

We then extracted the information of the layers for each plot and calculated the area occupied by each patch of habitat, which was successively converted to a percentage.

All spatial analyses were carried out using QGIS software. All the layers used were extracted from transportation network information from CNIG (<http://centrodedescargas.cnig.es/>). We reclassified the existing land cover classes in the land use layer (IGN; Cartografía Temática Ambiental of the Principado de Asturias, Hojas del Mapa de Vegetación, Litología, Roquedos y Hábitat del Oso. Escala 1:25000. © Principado de Asturias, Spain) into six habitat types: (1) deciduous forests; (2) conifer forests; (3) shrublands; (4) pastures; (5) rocky areas; and (6) crops.

Characterization of the study footpaths.—To analyse variability in the abundance of RTs along footpaths, we estimated the following variables for each footpath: 1) rub tree abundance, i.e., the kilometric abundance index of rub trees for each footpath (the ratio of the total number of RTs observed along a footpath with respect to the total footpath length covered at each site; hereafter, RT abundance on footpaths); (2) predominant RT species (categorical variable with 6 levels: birch, conifer, chestnut, oak, mixed, and other). Footpaths (length range 0.5 - 10.0 km) with at least 60% of RTs belonging to the same species were classified as paths characterized by that species, while footpaths without a predominant species were classified as ‘mixed’; (3) density of bear observations (see below) in a ring buffer of 1 km around each footpath where the RTs were found (hereafter, surrounding bear obs. density) as a proxy for the bear density around the RTs; and (4) the ratio of the distance to the nearest core cell with respect to the distance to the nearest border cell (ratio core/border). This last variable was a proxy for the relative position of the footpath in the area occupied by each population nucleus.

Explanatory variables for RT abundance.— We divided the area known to be

inhabited by the western subpopulation of brown bears in the Cantabrian Mountains into cells of 3×3 km that were classified into: 1) border cells, i.e., cells with at least 3 years of confirmed bear presence; 2) core area cells, i.e., cells with at least 7 years of confirmed bear presence; 3) core cells, i.e., cells within each core area that showed the highest number of years of confirmed bear presence. Cells not included in any of these categories were not considered to be inside the brown bear range and therefore were discarded (Zarzo-Arias et al. 2019; Fig.1). By undergoing this process, we obtained four different ‘core areas’ of bear distribution, i.e., cells with > 7 years of bear observations (Fig. 1), whereas the border of this distribution represented our border of the bear population. For each of these core areas, we selected one ‘core cell’, i.e., the cell with the most years of bear observations for each core area (Fig. 1). We calculated the distance from the central point of each footpath to: 1) the centroid of the nearest core cell (i.e., distance to core cell); and 2) the centroid of the nearest border cell (i.e., distance to border cell). We undertook this procedure in QGIS.

Brown bear observations used to obtain the values of the variables (i.e., surrounding density of bear observations, distance to core cell, and distance to border cell variables) were obtained for the period 2000 – 2017 by: (1) direct and indirect observations (scats, hairs and footprints) that were georeferenced by rangers of Principado de Asturias and Junta de Castilla y León, mainly by the ‘Patrulla Oso’ (Bear Patrol), as well as by all the other rangers of both regional governments, by the Asturian Foundation for the Conservation of Wildlife (FAPAS, Fondo para la Protección de los Animales Salvajes), by the Asturian Bear Foundation (FOA, Fundación Oso de Asturias) and the Brown Bear Foundation (FOP, Fundación Oso Pardo); (2) remotely triggered cameras that were randomly placed by the FAPAS and the Bear Patrol; and (3)

our own georeferenced observations (Sergiel et al. 2017; Lamamy et al. 2019; Penteriani et al. 2020).

Statistical analyses

In analysing the effect of individual tree characteristics on the probability of brown bear marking, it was necessary to account for the non-independence of rub and control trees sampled at the same location. We addressed this by adopting an approach that explicitly accounted for spatial autocorrelation using generalized additive models (GAMs; Hastie and Tibshirani 1986). This approach fits smoothing functions to easting and northing coordinates that account for non-linear spatial phenomena (i.e. spatial autocorrelation) and thus for the likelihood that coordinates close together will be more similar than those further apart. Given that rub and control trees have the same coordinates at a given sampling location, this approach controls for non-independence of these observations, as well as accounting for spatial autocorrelation at larger scales. GAMs were fitted with tree type as the binomial response variable (0 = control tree; 1 = rub tree) and the interaction between X and Y coordinates of trees as a smooth function (Wood 2004). We used the thin plate regression spline method with $k = 20$. This level of k was chosen through visual assessment of the residuals, to ensure sufficient smoothing while at the same time avoiding overfitting. To maintain relatively simple models for running the scenarios, and to minimise overfitting, smoothed terms were not used for the other continuous explanatory environmental variables (e.g. Gili et al. 2020). The following predictor variables were included in the models to be tested: tree species, tree height, trunk diameter, trunk height, tree spacing, and distance to footpath. All the variables were scaled, and there was no collinearity among explanatory variables (maximum value of Variance Inflation Factor, VIF = 2.7).

Next, we investigated whether RT abundance on footpaths (the kilometric abundance index) varied depending on local density of bear observations, location of the footpaths with respect to bear population distribution, and the predominant tree species around each footpath. Because RT abundance on footpaths was likely to vary between the two main portions of the bear population (i.e. Asturias and León), we first extracted the residuals of the linear model (LM) including the RT abundance on footpaths as the response variable and the geographic location (two-level variable: Asturias and León) as a predictor. These residuals thus included the variation derived from the location of each footpath (geographic location) in the final model. Because the residuals of the model did not follow a normal distribution, we carried out a logarithmic transformation of RT abundance on footpaths (Gelman and Hill 2006) to achieve normality. We then built LMs with these values as the response variable and the three above-mentioned parameters as explanatory variables.

We carried out all statistical analyses with the package “lme4” (Bates et al. 2015) and “mgcv” (Wood 2004) for GAMs in R v.3.5.2 statistical software (R Foundation for Statistical Computing 2018). To build both model classes (i.e., GAMs and LMs), we carried out the following procedure: to identify possible collinearity among predictor variables, we calculated variance inflation factors (VIF; Fox et al. 2007) for coefficients in the full model. Influence diagnostics were used to assess the presence of potential outliers and highly influential observations in our models, though none were found. We used model selection to test all combinations of the predictor variables (including the null model). Model selection was based on Akaike’s information criterion (AIC; Akaike 1973) using the library “MuMIn” (Bartoń 2013) and models with a $\Delta\text{AICc} < 2$ were considered as equally competitive. We then employed

multi-model averaging on the models with $\Delta AICc < 2$ to extract coefficients of each explanatory variable.

Ethical note

This article does not contain any studies with human participants or animals carried out by any of the authors. In this study, we used only observational data. For access to those areas restricted to the general public, we had the authorization from the pertinent environmental agencies of the regional governments of Castilla y León and the Principado de Asturias.

RESULTS

Location and characterisation of rub and control trees.—During this study, we found 101 RT and used 263 CT (Figure 1), with a range of 0 to 9 CT per RT. Only 3 of the RTs (<3%) and 8 of the CTs (3%) were dead. Rubbing marks such as scratches and bites were present in 81.2% and 80.2% of the RTs respectively. RTs were located at an average altitude of 895 m a.s.l. and were predominantly on north-facing slopes: 31.7% N ($n = 32$), 17.8% E ($n = 18$), 12.9% S ($n = 13$), 9.9% NE ($n = 10$), 9.9% NW ($n = 10$), 7.9% W ($n = 8$), 5.0% SE ($n = 5$) and SW ($n = 5$). Birches were the most frequently marked single tree type (28.7%), followed by oaks (16.3%), conifers (11.6%), and chestnuts (10.1%). Birches and conifers were marked at a higher percentage than that in which they appeared in the total pool of trees studied (18.6% and 7.5% respectively), while the rest of tree classes were marked in a similar proportion to their abundance. The results of the models testing for the effect of individual tree characteristics on the probability of being marked suggested that, among the variables considered, trunk DBH, mean distance to neighbouring trees, and tree species, were the most important

characters in explaining tree marking probability (Fig. 2; see also Supplementary Data SD1). Indeed, the best ranked model showed this combination of explanatory variables (Table 1). Specifically, RTs had larger DBHs and were in less dense parts of the forest stand (i.e., they showed higher distances from the nearest trees) compared to CTs. In addition, birches were the most frequently marked trees by bears among all tree classes (Tables 1 and 2). The smoothed spatial term was not significant in the full model ($\chi^2_2 = 2.36$, $P = 0.31$), and no model including this term was in the best model set (ΔAIC between the best model containing the smoothed term and the top model = 2.35), suggesting limited influence of spatial effects between RTs and CTs.

Characterization of the landscape around rub trees.—The landscape surrounding the RTs was covered predominantly by deciduous forest, followed by scrubland and pastures, without any human settlements within a 1 km radius (Supplementary Data SD2).

Characterization of the study footpaths.—To assess the kilometric abundance of RTs, we included 35 footpaths. These footpaths had 0 to 11 RTs, with an average of 3.7 RTs per footpath ($SD = 2.6$, $n = 35$). The mean RT abundance on footpaths was 2.5 RTs/km ($SD = 2.1$, $n = 35$). The footpaths located in areas with a predominance of birch showed on average a higher density of RTs (3.3 ± 3.5 RTs/km, $n = 9$) compared to footpaths dominated by conifers (2.4 ± 1.5 RTs/km, $n = 5$), chestnut trees (1.9 ± 0.4 RTs/km, $n = 3$), oaks (1.9 ± 0.9 RTs/km, $n = 4$), other species (3.0 ± 3.4 RTs/km, $n = 3$), and those without dominant species (mixed; 2.0 ± 0.7 RTs/km, $n = 11$). Along the footpaths, we detected some notably dense clusters of RTs, particularly in some footpaths located in birch forests (7 and 11 RTs in sections of 460 m and 930 m, respectively). In one case, we found a cluster of RTs in a homogeneous beech forest, with 7 RTs within 210 m of each other.

The set of models built to analyse potential factors affecting the intensity of bear rubbing, measured as the number of RTs along a footpath, showed that none of the variables under study played an important role. Indeed, the model with the lowest AICc value was the null model (Table 3). The second ranked model showed a weak positive relationship between RT abundance on footpaths and bear observation density (Table 3 and Fig. 3); however, the importance of this variable was very low (Tables 4).

DISCUSSION

In this study, we assessed brown bear selectivity of rub trees and the potential factors associated with marking behaviour along forest paths. It seems that Cantabrian brown bears select larger and more accessible trees, and that RT abundance is not determined by the density of bear observations or their geographical position within the bears' population range.

Specifically, brown bears preferentially used trees with greater trunk diameter. A preference for larger diameter trees also has been reported in studies spread throughout the brown bear range in North America (Green and Mattson 2003; Clapham et al. 2013), Russia (Puchkovskiy et al. 2012; Seryodkin 2014) and Japan (Sato et al. 2014). Trees with larger diameters may stand out over the surroundings and thus be more conspicuous (Green and Mattson 2003), which may also help if chemical signalling is associated with visual marks such as bites and scratches (Sato et al. 2014). This enhancement related to conspicuousness could be expected if the act of marking trees produces scent, as occurs when the bark is altered or when the marks are produced by pedal marking (Sergiel et al. 2017). Trunk height was almost significant, despite appearing in the best models, which probably suggests a trend towards greater height

between the ground and the first branches in the selected trees. Greater trunk heights (i.e., the available vertical space in which the bears can rub) have also been observed for rub trees in North America (Green and Mattson 2003; Clapham et al. 2013) and Russia (Seryodkin 2014), and might result in greater accessibility for signallers and facilitate the action of rubbing (Seryodkin 2014), particularly for the largest bears. Rub trees were predominantly located on north-facing slopes, as also reported by Seryodkin (2014). Tattoni et al. (2015), in a study with camera traps, detected a higher cumulative camera trapping rate in NE and S aspects, orientations that in this study represented the 3rd and 4th position. Because most of the tree species included in this study develop better in the wetter conditions of northern slopes, e.g., beeches (Sánchez et al. 2003), chestnuts (Blanco Andray et al. 2000), sessile oaks (Díaz-Maroto et al. 2006), and birches (García et al. 2005), it seems plausible that the greater abundance of brown bear RTs is related not to the exposure itself, but to the fact that on these northern slopes forest stands are better developed and the trees grow larger than on the southern slopes.

Although our RT surveys were carried out exclusively along footpaths, we highlight here that in most studies on bear marking, trees were predominantly located along routes, game trails, and forest edges. Indeed, increasing distance to the footpath represented an explanatory, negatively-related variable with the probability of rubbing in other studies (86% in Green and Mattson 2003 and 100% in Clapham et al. 2013). Similarly, higher intensity of usage of RTs on large trails and forestry roads have been described in the Italian Alps, probably because they are preferred movement routes by bears (Tattoni et al. 2015). Some authors argue that observations of rubbing hairs, bites and scratches oriented towards roads and footpaths represent strong evidence that bears prefer to perform chemical signalling specifically on those trees located on the sides of footpaths and trails (Lloyd 1979; Burst and Pelton 1983; Green and Mattson 2003; Nie

et al. 2012), which also are routes of brown bear displacement (Burst and Pelton 1983; Seryodkin 2014).

Even though we did not carry out an extensive census of all the tree species within the marking tree stand, we established whether the RT species were similar to those in their surroundings by comparing the RTs with their surrounding CTs. Thus, we discovered that birch was the tree species most frequently marked by brown bears, because it was found in greater proportion among marked trees than among all trees. Conifers also were marked in a greater proportion to their availability. However, the conifer species marked by bears in our study (Monterey pine [*Pinus radiata*] and Douglas fir [*Pseudotsuga menziesii*]) do not occur naturally in the study area, and the individuals included corresponded to plantations or groups of feral trees. Conifers are frequently identified as preferred rub trees because of the aromas they produce when they are lacerated (Puchkovskiy 2009; Nie et al. 2012; Clapham et al. 2013; Sato et al. 2014), which can enhance or help maintain the bear's scent and/or attract recipient bears. However, our study area did not have enough locations where coniferous and deciduous species co-occur to test for a potential preference for conifer rubbing by Cantabrian brown bears.

In our study area, birch is distributed irregularly, generally in acidic and wet soils or near water at forest boundaries or in cleared areas, forming continuous forests only in a few areas at high elevation (García de Celis et al. 2004). Nevertheless, it was proportionally the most marked tree species. For instance, in those places where several birches occurred along a path, most of them were marked, sometimes every few meters. Similar findings have been reported by studies conducted in boreal conifer-dominated forests, where birches were found to be marked more frequently than other deciduous species. For example, Puchkovskiy (2009) reported that in several forests of the Russian

taiga, birch was the most frequently marked deciduous tree and that when birches were present, brown bears always chose it preferentially over other deciduous species. In addition, Seryodkin (2014) described the appearance of brown bear marking trees in Kamchatka forests dominated by the stone birch *Betula ermanii*. Preference for birch species also was reported in the Middle Sikhote-Alin, SE Russia (Seryodkin et al. 2014), suggesting that brown bears selected birch not only for their availability, but also for physiognomic characteristics that favour marking. In fact, birches tend to have a large diameter with no branches on the first few metres of the trunk, and the stratified bark is lacerated easily by scratches and bites. Also, the birch has a whitish outer layer contrasting with a dark inner layer, so when it is removed or lacerated the marks are very evident, even after healing, which may increase the conspicuousness of brown bear visual markings. In addition, in the common case of a birch RT surrounded by other tree species, generally characterised by darker barks (*Quercus* spp., beech, chestnut, etc.), the RT itself would stand out in the surroundings, thus making its visual traceability easier for bears and therefore becoming an advantageous and long-lasting substrate for marking. Marking has been suggested to include visual and olfactory signalling in different bear species, including brown bears (e.g., Sergiel et al. 2017), and different types of marking behavior have been documented in social and solitary carnivores (e.g., Paquet 2011; Vogt et al. 2014).

It is worth noting that the sections dominated by birches had the highest rubbing densities. RT densities in the Cantabrian Mountains are higher than those recorded by Henderson et al. (2015) in conifer forests of the northwest US, where they compared RT abundances between developed trails and roads (1.0 ± 1.1 RTs/km; $n = 30$) and game trails (0.8 ± 1.1 RTs/km; $n = 30$). In several Kamchatka valleys dominated by stone birch, Seryodkin (2014) reported very variable abundances (from 40 RTs / km to 0.4

RTs / km) and RTs groupings in short sections of the footpath (7 RTs in 20 m or 5 RTs in 8 m). We detected similar but lower density aggregations, especially in birch-dominated forests. This indicates a trend towards more intense or greater marking on them, supporting the positive selectivity towards birches described above. Understanding wildlife behaviour can prove useful for conservation and management (e.g., Greggor et al. 2019). In the context of our study, the noticeable selection of birch by marking bears might provide a tool to monitor bear presence, e.g., in areas immediately surrounding the present range of the species, where targeting birches in monitoring programs that search for bear signs might help detect dispersing individuals.

To our knowledge, this is the first time that the density of brown bear rub trees has been compared with some features of the bear population or its proxies. In our study, neither the density of bear observations (as a proxy for bear density), nor the position of the track within the bear distribution area, affected the kilometric abundance of RTs. Because the null model had the lowest AIC, we deduce that the determinants of the intensity of rubbing in our population are different from those analysed in this study, hence the assessment of other possible variables may be necessary. The lack of an effect of the density of observations on the density of rub trees may be due to the fact that differences in bear density, or more specifically of bear observations around each foot path, are not large enough to affect abundance from rub trees. More research in this field is needed to elucidate whether variations in bear density can affect rubbing behaviour in other ways, such as modifications in the frequency of rubbing or the proportion of rubbing between sex and age classes, as Lamb et al. suggested (2017). In one study considering the distribution of rubbing sites in a territorial carnivore, the tiger (*Panthera tigris*), rubbing increased towards the limits of the territory (Smith et al. 1989). This might not be applicable to brown bears that are non-territorial species in

which individuals of the same population can have overlapping home ranges (Seryodkin et al. 2017, Frank et al. 2018).

There are three potential biases in this study. First, searches for RTs were undertaken mostly (but not exclusively) from footpaths. The findings that RTs were closer to, and orientated towards, footpaths may thus have been the result of greater detectability from those paths. However, we do not believe this to have affected our results given the large amount of evidence that footpaths are common routes of brown bear displacement and that RTs occur more frequently along such routes (e.g., Nie et al. 2012; Seryodkin 2014; see above). Second, the lighter coloured bark of birch trees may have made them more conspicuous to observers. We believe this potential bias was minimised by very careful observations of all potential RTs. Furthermore, preference for birch has been found in several other studies (see above). Third, we used proxy measures to estimate bear density and distribution. We cannot know whether our measures are fully accurate representations of the population but nevertheless, our measures were the best available, being based largely on observations made by trained forest rangers and researchers. While we believe the above potential biases were minimised in our study, their effects only can be properly elucidated through further research based on intensive telemetry studies.

To conclude, our results suggest that Cantabrian brown bears select rub trees based on characteristics related to the tree, rather than their position relative to areas with a high density of conspecifics and/or with the distribution of individuals within the population range. Thus, even in small and isolated bear populations, such as the one under study here, some aspects of rubbing behaviour such as tree selection and marking effort may not be affected by population size and/or structure.

528

529

ACKNOWLEDGEMENTS

530 We thank the Administrations of the Government of the Principality of Asturias (in
531 particular, we thank Teresa Sánchez, Víctor Vázquez, and Pedro García-Rovés) and the
532 Junta de Castilla y León (especially David Cubero, Mercedes García, and M. Ángeles
533 Osorio) for providing the bear database and for their help during this study. We also
534 thank M. Fernández, D. Ramos, and F. Somoano, of the ‘Patrulla Oso’ (Bear Patrol) of
535 Asturias for their continuous help in the field.

536

FUNDING

537 During this research, EGB was financially supported by a FPU grant (FPU15-03429) of
538 the Spanish Ministry of Science, Innovation, and Universities. V. P. and A. O. were
539 financially supported by the Excellence Project CGL2017-82782-P financed by the
540 Spanish Ministry of Science, Innovation, and Universities, the Agencia Estatal de
541 Investigación (AEI) and the Fondo Europeo de Desarrollo Regional (FEDER, EU). V.
542 P. also was financially supported by a GRUPIN research grant from the Regional
543 Government of Asturias (Ref.: IDI/2018/000151). G. B. was financially supported by a
544 collaboration contract with the MUSE – Museo delle Scienze of Trento (Italy).

545

546

SUPPLEMENTARY DATA

547 **Supplementary Data SD1.** — Mean value, standard deviation (SD) and minimum (Min)
548 and maximum (Max) values of each of the parameters used to characterise both rub and
549 control trees.

Supplementary Data SD2. — Mean value, standard deviation (SD) and range of landscape features surrounding rub trees, including the- percent of the different land cover within a 1km buffer around each rub tree.

LITERATURE CITED

ALBERTS, A. C. 1992. Constraints on the design of chemical communication systems in terrestrial vertebrates. *American Naturalist* 139:S62–S89.

ALLEN, M. L., L. HOČEVAR, M. DE GROOT, AND M. KROFEL. 2017. Where to leave a message? The selection and adaptive significance of scent marking sites for Eurasian lynx. *Behavioral Ecology and Sociobiology* 71:136–145.

BARJA, I. 2009. Decision making in plant selection during the faecal-marking behaviour of wild wolves. *Animal Behaviour* 77:489–493.

BARJA, I., AND F. J. DE MIGUEL. 2010. Chemical communication in large carnivores: Urine-marking frequencies in captive tigers and lions. *Polish Journal of Ecology* 58:397–400.

BARTOŃ, K. 2013. Model selection and model averaging based on information criteria (AICc and alike). *The Comprehensive R Archive Network* 1:13.

BATES, D., M. MAECHLER, B. BOLKER, AND S. WALKER. 2015. Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software* 67:1–48.

BLANCO ANDRAY, A., A. RUBIO SÁNCHEZ, O. SÁNCHEZ PALOMARES, R. ELENA ROSSELLÓ, V. GÓMEZ SANZ, AND S. GRAÑA DOMÍNGUEZ. 2000. Autoecología de los castaños de Galicia (España). *Investigación Agraria. Sistemas y Recursos Forestales* 9:337–360.

573 BROMLEY, G. F. 1965. Bears of the south of the USSR Far East. Nauka press, Moscow-
574 Leningrad, Russia.

575 BROWN, R. E. 1979. Mammalian social odors: a critical review. *Advances in the Study*
576 *of Behavior* 10:103–162.

577 BURST, T. L., AND M. R. PELTON. 1983. Black bear mark trees in the Smoky Mountains.
578 *Bears: Their Biology and Management* 5:45–53.

579 CLAPHAM, M., O. T. NEVIN, A. D. RAMSEY, AND F. ROSELL. 2012. A hypothetico-
580 deductive approach to assessing the social function of chemical signalling in a non-
581 territorial solitary carnivore. *PLoS ONE* 7:1–11.

582 CLAPHAM, M., O. T. NEVIN, A. D. RAMSEY, AND F. ROSELL. 2013. The function of
583 strategic tree selectivity in the chemical signalling of brown bears. *Animal*
584 *Behaviour* 85:1351–1357.

585 CLAPHAM, M., O. T. NEVIN, A. D. RAMSEY, AND F. ROSELL. 2014. Scent-marking
586 investment and motor patterns are affected by the age and sex of wild brown bears.
587 *Animal Behaviour* 94:107–116.

588 CLAPPERTON, B. K. 1989. Scent-marking behaviour of the ferret, *Mustela furo* L.
589 *Animal Behaviour* 38:436–446.

590 CLEVINGER, A. P., AND F. J. PURROY. 1991. Demografía del oso pardo (*Ursus arctos*)
591 en la Cordillera Cantábrica. *Ecología* 5:243–256.

592 DÍAZ-MAROTO, I. J., P. VILA-LAMEIRO, AND M. C. DÍAZ-MAROTO. 2006. Autecology of
593 sessile oak (*Quercus petraea*) in the north-west Iberian Peninsula. *Scandinavian*
594 *Journal of Forest Research* 21:458–469.

595 DOTY, R. L. 1986. Odor-guided behavior in mammals. *Experientia* 42:257–271.

596 FERNÁNDEZ-GIL, A., J. NAVES, AND M. DELIBES. 2006. Courtship of brown bears *Ursus*
597 *arctos* in northern Spain: phenology, weather, habitat and durable mating areas.
598 *Wildlife Biology* 12:367–373.

599 FILIPCZYKOVÁ, E., I. M. A. HEITKÖNIG, A. CASTELLANOS, W. HANTSON, AND S. M. J. G.
600 STEYAERT. 2016. Marking behavior of Andean bears in an Ecuadorian cloud
601 forest: a pilot study. *Ursus* 27:122–128.

602 FOX, J., ET AL. 2007. The “car” package. R Foundation for Statistical Computing.

603 FRANK, S. C., ET AL. 2018. Sociodemographic factors modulate the spatial response of
604 brown bears to vacancies created by hunting. *Journal of Animal Ecology* 87:247–
605 258.

606 GARCÍA-GARITAGOITIA, J. L., I. R. FRAILE, AND I. D. VILLAREJO. 2007. Variabilidad
607 genética. Pages 69–90 in *Demografía, distribución, genética y conservación del*
608 *oso pardo cantábrico* (G. Palomero, F. Ballesteros, J. Herrero, and C. Nores, eds.).
609 *Naturaleza y Parques Nacionales, Serie Técnica, Ministerio de Medio Ambiente.*
610 *Madrid, Spain.*

611 GARCÍA, D., M. QUEVEDO, J. R. OBESO, AND A. ABAJO. 2005. Fragmentation patterns
612 and protection of montane forest in the Cantabrian range (NW Spain). *Forest*
613 *Ecology and Management* 208:29–43.

614 GARCÍA DE CELIS, A. J., J. C. GUERRA VELASCO, AND L. C. MARTÍNEZ FERNÁNDEZ.
615 2004. Los abedulares de la Omaña Alta (León): notas sobre dinámica vegetal y
616 cambios en el aprovechamiento de los montes. *Boletín de la A.G.E* 38:245–258.

617 GILL, F., S. E. NEWSON, S. GILLINGS, D. E. CHAMBERLAIN, AND J. A. BORDER. 2020. Bats
618 in urbanising landscapes: habitat selection and recommendations for a sustainable

619 future. *Biological Conservation* 241:108343.

620 GONZALEZ, E. G., J. C. BLANCO, F. BALLESTEROS, L. ALCARAZ, G. PALOMERO, AND I.
621 DOADRIO. 2016. Genetic and demographic recovery of an isolated population of
622 brown bear *Ursus arctos* L., 1758. *PeerJ* 4:1–25.

623 GORMAN, M. L., AND B. J. TROWBRIDGE. 1989. The role of odor in the social lives of
624 carnivores. Pages 57–88 in *Carnivore behavior, ecology, and evolution* (J. L.
625 Gittleman, ed.). Cornell University Press. Ithaca, New York.

626 GOSLING, L. M. 1990. Scent marking by resource holders: alternative mechanisms for
627 advertising the costs of competition. Pages 315–328 in *Chemical signals in*
628 *vertebrates 5* (D. W. Müller-Schwarze and S. E. Natynczuk, eds.). Oxford
629 University Press. Oxford, United Kingdom.

630 GOSLING, L. M., AND H. V. MCKAY. 1990. Competitor assessment by scent matching:
631 an experimental test. *Behavioral Ecology and Sociobiology* 26:415–420.

632 GOSLING, L. M., AND S. C. ROBERTS. 2001. Scent-marking by male mammals: cheat-
633 proof signals to competitors and mates. Pages 169–217 in *Advances in the Study*
634 *of Behavior* (P. J. B. Slater, J. S. Rosenblatt, C. T. Snowdon, and T. J. Roper, eds.).
635 Academic Press. San Diego, California.

636 GREEN, G. I., AND D. J. MATTSON. 2003. Tree rubbing by Yellowstone grizzly bears
637 *Ursus arctos*. *Wildlife Biology* 9:1–9.

638 GREGGOR, A. L., D. T. BLUMSTEIN, B. B. M. WONG, AND O. BERGER-TAL. 2019. Using
639 animal behavior in conservation management: A series of systematic reviews and
640 maps. *Environmental Evidence* 8:1–3.

641 HASTIE, T., AND R. TIBSHIRANI. 1986. Generalized additive models. *Statistical science*

642 1:297–318.

643 JOHNSTON, R. E. 2008. Individual odors and social communication: individual
644 recognition, kin recognition, and scent over-marking. *Advances in the Study of*
645 *Behavior* 38:439-505.

646 JOHNSTON, R. E., AND J. DEL BARCO-TRILLO. 2010. Communication by chemical
647 signals: Behavior, social recognition, hormones and the role of the vomeronasal
648 and olfactory systems. *Hormones, Brain and Behavior Online*:395–441.

649 JOJOLA, S. M., F. ROSELL, I. WARRINGTON, J. E. SWENSON, AND A. ZEDROSSER. 2012.
650 Subadult brown bears (*Ursus arctos*) discriminate between unfamiliar adult male
651 and female anal gland secretion. *Mammalian Biology* 77:363–368.

652 KLEIMAN, D. 1966. Scent marking in the Canidae. *Symposia of the Zoological Society*
653 *of London* 18:167–177.

654 KLEINER, J. D., R. C. VAN HORN, J. E. SWENSON, AND S. M. J. G. STEYAERT. 2018. Rub-
655 tree selection by Andean bears in the Peruvian dry forest. *Ursus* 29:58–66.

656 KROTT, P. 1962. Beiträge zur Kenntnis des Alpenbärs, *Ursus arctos* Linné 1758.
657 *Säugetierkundliche Mitteilungen* 10:1–35. [In German].

658 LAMAMY, C., G. BOMBIERI, A. ZARZO-ARIAS, E. GONZÁLEZ-BERNARDO, AND V.
659 PENTERIANI. 2019. Can landscape characteristics help explain the different trends
660 of Cantabrian brown bear subpopulations? *Mammal Research*:1–9.

661 LAMB, C. T., G. MOWAT, S. L. GILBERT, B. N. McLELLAN, S. E. NIELSEN, AND S.
662 BOUTIN. 2017. Density-dependent signaling: An alternative hypothesis on the
663 function of chemical signaling in a non-territorial solitary carnivore. *PLoS ONE*
664 12:1–12.

665 LAURIE, A., AND J. SEIDENSTICKER. 1977. Behavioural ecology of the Sloth bear
666 (*Melursus ursinus*). Journal of Zoology 182:187–204.

667 LLOYD, K. A. 1979. Aspects of the ecology of black and grizzly bears in coastal British
668 Columbia. MS thesis, University of British Columbia. Vancouver, British
669 Columbia, Canada.

670 MACDONALD, D. W. 1980. Patterns of scent marking with uring and faeces amongst
671 carnivore communities. Symposia of the Zoological Society of London 45:107–
672 139.

673 MARTÍNEZ CANO, I., F. G. TABOADA, J. NAVES, A. FERNÁNDEZ-GIL, AND T. WIEGAND.
674 2016. Decline and recovery of a large carnivore: environmental change and long-
675 term trends in an endangered brown bear population. Proceedings of the Royal
676 Society B: Biological Sciences 283:20161832.

677 MATEO-SÁNCHEZ, M. C., ET AL. 2016. Seasonal and temporal changes in species use of
678 the landscape: how do they impact the inferences from multi-scale habitat
679 modeling? Landscape Ecology 31:1261–1276.

680 MILLS, M. G. L., M. L. GORMAN, AND M. E. J. MILLS. 1980. The scent marking
681 behaviour of the brown hyaena *Hyaena brunnea*. South African Journal of
682 Zoology 15:240–248.

683 MORALES-GONZÁLEZ, A., H. RUÍZ-VILLAR, A. OZGUL, N. MAAG, AND G. COZZI. 2019.
684 Group size and social status affect scent marking in dispersing female meerkats.
685 Behavioral Ecology 30:1602–1610.

686 MORGAN HENDERSON, M. J., M. HEBBLEWHITE, M. S. MITCHELL, J. B. STETZ, K. C.
687 KENDALL, AND R. T. CARLSON. 2015. Modeling multi-scale resource selection for

688 bear rubs in northwestern Montana. *Ursus* 26:28–39.

689 MÜLLER-SCHWARZE, D. 2006. Chemical ecology of vertebrates. Cambridge University
690 Press. Cambridge, United Kingdom.

691 MÜLLER, C. A., AND M. B. MANSER. 2008. Scent-marking and intrasexual competition
692 in a cooperative carnivore with low reproductive skew. *Ethology* 114:174–185.

693 NASA. 2019. NASA GLOBE Observer App (v 3.0).

694 NAVES, J., AND G. PALOMERO. 1993. Distribución del oso pardo en la Cordillera
695 Cantábrica. Pages 35–46 in *El oso pardo (Ursus arctos) en España* (J. Naves and
696 G. Palomero, eds.). ICONA Colección Técnica. Madrid, Spain.

697 NIE, Y., R. R. SWAISGOOD, Z. ZHANG, Y. HU, Y. MA, AND F. WEI. 2012. Giant panda
698 scent-marking strategies in the wild: Role of season, sex and marking surface.
699 *Animal Behaviour* 84:39–44.

700 ORTEGA, M. T., AND C. G. MORALES. 2015. El clima de la Cordillera Cantábrica
701 castellano-leonesa: diversidad, contrastes y cambios [The climate of the Castilian-
702 Leonese Cantabrian Mountains: diversity, contrasts and changes]. *Investigaciones*
703 *Geográficas* 63:45–67.

704 PATO, J., AND J. R. OBESO. 2012. Growth and reproductive performance in bilberry
705 (*Vaccinium myrtillus*) along an elevation gradient. *Ecoscience* 19:59–68.

706 PENTERIANI, V., AND M. DELGADO. 2017. Living in the dark does not mean a blind life:
707 Bird and mammal visual communication in dim light. *Philosophical Transactions*
708 *of the Royal Society B: Biological Sciences* 372:1–8.

709 PENTERIANI, V., ET AL. 2020. Females brown bear use areas with infanticide risk in a
710 spatially confined population. *Ursus* 31:1–9.

711 PENTERIANI, V., A. ZARZO-ARIAS, A. NOVO-FERNÁNDEZ, G. BOMBIERI, AND C. A.
 712 LÓPEZ-SÁNCHEZ. 2019. Responses of an endangered brown bear population to
 713 climate change based on predictable food resource and shelter alterations. *Global*
 714 *Change Biology* 25:1133–1151.

715 PÉREZ, T., ET AL. 2014. Estimating the population size of the endangered Cantabrian
 716 brown bear through genetic sampling. *Wildlife Biology* 20:300–309.

717 PÉREZ, T., ET AL. 2010. Evidence for improved connectivity between Cantabrian brown
 718 bear subpopulations. *Ursus* 21:104–108.

719 PUCHKOVSKIY, S. V. 2009. Selectivity of tree species as activity target of brown bear in
 720 taiga. *Contemporary Problems of Ecology* 2:260–268.

721 PUCHKOVSKIY, S. V., M. S. BUINOVSKAYA, D. K. VORONETSKAYA, AND G. V.
 722 NEUSTROEV. 2012. On the studies of marking behavior of brown bear in terms of
 723 tree diameter selectivity. *Contemporary Problems of Ecology* 5:104–109.

724 R Foundation for Statistical Computing. 2018. R: A Language and Environment for
 725 Statistical Computing. Vienna, Austria.

726 RALLS, K. 1971. Mammalian scent marking. *Science* 171:443–449.

727 RICH, T. J., AND J. L. HURST. 1998. Scent marks as a reliable indicator of the
 728 competitive ability of mates. *Animal Behaviour* 56:727–735.

729 Roberts, S. C., and L. M. Gosling. 2001. The economic consequences of advertising
 730 scent mark location on territories. Pages 11–17 *Chemical signals in vertebrates 9*
 731 (A. Marchlewska-Koj, J. J. Lepri, and D. Müller-Schwarze, eds.). Springer US.
 732 Boston, Massachusetts.

733 ROSELL, F., ET AL. 2011. Brown bears possess anal sacs and secretions may code for

734 sex. Journal of Zoology 283:143–152.

735 SÁNCHEZ, O., A. RUBIO, A. BLANCO, R. ELENA, AND V. GÓMEZ. 2003. Autoecología
736 paramétrica de los hayedos de Castilla y León. Investigación Agraria. Sistemas y
737 Recursos Forestales 12:87–110.

738 SATO, Y., ET AL. 2014. Selection of rub trees by brown bears (*Ursus arctos*) in
739 Hokkaido, Japan. Acta Theriologica 59:129–137.

740 SAWAYA, M. A., J. B. STETZ, A. P. CLEVINGER, M. L. GIBEAU, AND S. T. KALINOWSKI.
741 2012. Estimating grizzly and black bear population abundance and trend in banff
742 national park using noninvasive genetic sampling. PLoS ONE 7:1–12.

743 SERGIEL, A., ET AL. 2017. Histological, chemical and behavioural evidence of pedal
744 communication in brown bears. Scientific Reports 7:1052.

745 SERYODKIN, I. V. 2014. Marking activity of the Kamchatka brown bear (*Ursus arctos*
746 *piscator*). Achievements in the Life Sciences 8:153–161.

747 SERYODKIN, I. V., A. V. KOSTYRIA, AND J. M. GOODRICH. 2014. Marking activities of
748 the brown bear in the Sikhote-Alin Mountains. Russian Journal of Zoology
749 95:694–702.

750 SERYODKIN, I. V, J. PACZKOWSKI, M. Y. BORISOV, AND Y. K. PETRUNENKO. 2017. Home
751 ranges of brown bears on the Kamchatka Peninsula and Sakhalin Island.
752 Contemporary Problems of Ecology 10:599–611.

753 SHAFFER, S. C. 1971. Some biological relationships of grizzly bears and black bears of
754 the Apgar Mountains in Glacier National Park, Montana. MS thesis, University of
755 Montana. Missoula, Montana.

756 SMITH, J. L. D., C. MCDUGAL, AND D. MIQUELLE. 1989. Scent marking in free-ranging

757 tigers, *Panthera tigris*. *Animal Behaviour* 37:11–27.

758 SWAISGOOD, R. R., D. G. LINDBURG, A. M. WHITE, H. ZHANG, AND X. ZHOU. 2004.

759 Chemical communication in giant pandas; experimentation and application. Pp.

760 106–119 in *Giant pandas: biology and conservation* (D. Lindburg and K.

761 Baragona, eds.). University of California Press. Berkeley, California.

762 TATTONI, C., N. BRAGALANTI, C. GROFF, AND F. ROVERO. 2015. Patterns in the use of

763 rub trees by the Eurasian brown bear. *Hystrix* 26:118–124.

764 TOMIYASU, J., ET AL. 2017. Testosterone-related and seasonal changes in sebaceous

765 glands in the back skin of adult male brown bears (*Ursus arctos*). *Canadian Journal*

766 *of Zoology* 96:205–211.

767 WHITE, A. M., R. R. SWAISGOOD, AND H. ZHANG. 2002. The highs and lows of chemical

768 communication in giant pandas (*Ailuropoda melanoleuca*): Effect of scent

769 deposition height on signal discrimination. *Behavioral Ecology and Sociobiology*

770 51:519–529.

771 WOOD, S. N. 2004. Stable and efficient multiple smoothing parameter estimation for

772 generalized additive models. *Journal of the American Statistical Association*

773 99:673–686.

774 ZARZO-ARIAS, A., ET AL. 2018. Brown bear behaviour in human-modified landscapes :

775 The case of the endangered Cantabrian population, NW Spain. *Global Ecology and*

776 *Conservation* 16:1-14.

777 ZARZO-ARIAS, A., ET AL. 2019. Identifying potential areas of expansion for the

778 endangered brown bear (*Ursus arctos*) population in the Cantabrian Mountains

779 (NW Spain). *PLoS ONE* 14:1–15.

FIGURE LEGENDS

Fig. 1.—Locations of the RTs included in the study and the range of the western subpopulation of Cantabrian brown bear, divided into cells of different observation frequency. Inset shows location of mapped area within the geographic confines of Spain. Locations of the 101 brown bear rub trees located in the Cantabrian Mountains (NW Spain; provinces of Asturias and León) are represented with purple dots. Due to the short distance that separates rub trees in some cases, several trees can be under the same spot. The range of the western subpopulation of Cantabrian brown bear was calculated as 3x3 km cells with at least three continuous years of bear observations (since 2000; Lamamy et al. 2019). The cells with only three continuous years of bear observations are in blue and were considered as the ‘border cells’. The core areas, i.e., areas with at least seven years of bear presence, are in orange. Within the core areas, we calculated the ‘core cells’ (in red), i.e., cells within each of the core areas that showed the highest number of continuous years of bear presence.

Fig. 2.—Characteristics of parameters used to describe rub trees (RT) used by brown bears ($n = 101$ RT; dark grey) and control trees (CT; $n = 363$; light grey) in the Cantabrian Mountains of NW Spain.

Fig. 3.— Changes of the logarithm of the RT abundance on footpaths of brown bear rub trees (LogIKA) along footpaths in the Cantabrian Mountains with respect to a proxy of brown bear density (left panel) and the position of the trail (right panel) within the bear population’s range (see also Fig. 2). Details on the proxy of bear density and the ratio core/border distance are provided in the main text.

TABLES

Table 1.—Competing models from GAMs built to investigate individual tree characteristics ($n = 364$ trees) determining marking probability in the Cantabrian brown bear population. Competitive models are ranked from the lowest (best model) to the highest AICc value. Only models with $\Delta\text{AICc} < 2$ are shown. Binomial response variable: tree type (0 = control tree (CT) and 1 = marked tree (RT)). R-squared of the most parsimonious model is 0.26. Note that a smoothed spatial coordinate term was included in the full model, but it was not selected in any of the top models.

Dependent variable	Competing models	df	AICc	ΔAICc	Weight
<i>Tree type</i> (CT/RT)	Trunk diameter + tree spacing + trunk height + tree species	8	368.58	0.00	0.23
	Trunk diameter + tree spacing + tree height + trunk height + tree species	9	368.91	0.32	0.20
	Trunk diameter + tree spacing + tree species	7	369.30	0.72	0.16
	dist. to footpath + trunk diameter + tree spacing + tree height + trunk height + tree species	10	369.36	0.78	0.16
	dist. to footpath + trunk diameter + tree spacing + trunk height + tree species	9	369.46	0.88	0.15
	Distance to nearest footpath + trunk diameter + tree spacing + tree species	8	370.18	1.60	0.10

Table 2.—Effects of individual tree characteristics ($n = 364$ trees) on the probability that a given tree was a bear rubbing tree in the Cantabrian Mountains. For each explanatory variable, we report the estimate (β), standard error (SE), significance (p), confidence intervals (CI) and relative importance values (RIV) obtained from model averaging on the models with $\Delta AICc < 2$. Binomial response variable: tree type (0 = control tree (CT) and 1 = marked tree (RT)). Baseline level for the categorical variable tree species is “other”. The most important parameters are highlighted in bold.

Dependent variable	Explanatory variable	Model-averaged coefficients and relative importance values				
		β	SE	p	CI	RIV
<i>Tree type</i> (CT/RT)	Intercept	-1.088	0.230	2.40e-06	(-1.539; -0.637)	-
	Trunk diameter	0.917	0.226	5.14e-05	(0.474; 1.360)	1.00
	Tree spacing	0.698	0.143	1.10e-06	(0.419; 0.979)	1.00
	Trunk height	0.275	0.152	0.072	(-0.023; 0.573)	0.73
	Birch	0.834	0.385	0.031	(0.079; 1.589)	1.00
	Chestnut	-0.363	0.447	0.418	(-1.239; 0.513)	1.00
	Conifer	0.055	0.602	0.928	(-1.125; 1.235)	1.00
	Oak	-0.558	0.383	0.146	(-1.309; 0.193)	1.00
	Tree height	-0.301	0.217	0.168	(-0.726; 0.124)	0.35
	Dist. to footpath	-0.181	0.167	0.279	(-0.508; 0.093)	0.41

Table 3.—Competing models built to investigate the effect of ecological characteristics of the surroundings of the study footpaths ($n = 35$) on the abundance of brown bear rub trees. Competitive linear models are ranked from the lowest (best model) to the highest AICc value. Only models with $\Delta\text{AICc} < 2$ are shown. R-squared of the model including bear obs. density = 0.062.

Dependent variable	Competing models	df	AICc	ΔAICc	Weight
RT abundance on footpath residuals	Null model	2	66.50	0.00	0.52
	Surrounding bear obs. density	3	66.66	0.16	0.48

Table 4.—Effects of ecological characteristics of the surroundings of the footpaths ($n = 35$) located in the Cantabrian Mountains on the abundance of brown bear rub trees. For each explanatory variable, we report the estimate (β), standard error (SE), significance (p), confidence intervals (CI), and relative importance values (RIV) obtained from model averaging on the models with $\Delta AICc < 2$.

Dependent variable	Explanatory variable	Model-averaged coefficients and relative importance values				
		β	SE	p	CI	RIV
RT abundance on footpath residuals	Intercept	3.354e-18	9.998e-02	1.000	(-0.196; 0.196)	-
	Bear obs. density	1.485e-01	1.005e-01	0.155	(-0.048; 0.345)	0.48